

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets

(11) Publication number:

0 053 434
A1

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 81305128.1

(51) Int. Cl.³: **C 07 C 101/18, C 07 C 101/62,**
C 07 C 101/00

(22) Date of filing: 29.10.81

(30) Priority: 28.11.80 US 211340

(71) Applicant: **AMERICAN HOSPITAL SUPPLY CORPORATION**, One American Plaza, Evanston, IL 60201 (US)

(43) Date of publication of application: **09.06.82**
Bulletin 82/23


(72) Inventor: **Erhardt, Paul W.**, 811 Cambridge Drive, Mundelein Illinois 60060 (US)
Inventor: **Borgman, Robert J.**, 1735 Victoria Road, Mundelein Illinois 60060 (US)
Inventor: **O'Donnell, John P.**, 109 Williams Street, Morgantown West Virginia 26505 (US)

(84) Designated Contracting States: **AT BE CH CH DE FR GB IT LU SE**

(74) Representative: **Seaborn, George Stephen et al**, c/o Edward Evans & Co. Chancery House 53-64 Chancery Lane, London WC2A 1SD (GB)

(54) **Compounds and method for treatment or prophylaxis of cardiac disorders.**

(57) A short-acting β -blocking compound of the formula



wherein AR may be substituted or unsubstituted aromatic, Y may be a straight or branched carbon chain or aralkyl, R may be lower alkyl, lower alkenyl, lower alkynyl, aryl or aralkyl, and x is an integer from 1 to about 3; or a pharmaceutically acceptable salt thereof.

EP 0 053 434 A1

COMPOUNDS AND METHOD FOR TREATMENT OR
PROPHYLAXIS OF CARDIAC DISORDERS

Background of the Invention

5 The present invention relates to the treatment or prophylaxis of cardiac disorders. More particularly, the invention relates to a novel method of treatment or prophylaxis of cardiac disorders which comprises administration of β -adrenergic blocking agents and to compounds useful in such method.

10

 The therapeutic and prophylactic uses of compounds which block sympathetic nervous stimulation of β -adrenergic receptors in the heart, lungs, vascular system and other organs are well documented. Typically, such compounds are administered therapeutically to patients
15 suffering from ischemic heart disease or myocardial infarction for the purpose of reducing heart work, i.e., heart rate and contractile force. Reducing heart work reduces oxygen demand, and may also actually increase oxygen supply. Thus reducing heart work can aid in the prevention of further tissue damage and can relieve angina pectoris.

20

β -Adrenergic stimulation may also aggravate or cause arrhythmias because of increased levels of catecholamines. Thus β -blocking agents may be employed to reduce the risks of arrhythmias.

25


 Compounds have been discovered which selectively block β -adrenergic receptors in various organs. Beta receptors in the heart are generally referred to as β_1 receptors, and those associated with vasodilation and bronchodilation are β_2 receptors. Non-selective β -blockers are not preferred for the treatment of cardiac disorders
30 because of their hypertensive action and potentiation undesirable effects on asthmatic patients. A number of β_1 selective adrenergic blocking

agents have been discovered. Smith, L.H., J. Appl. Chem. Biotechnol., 28, 201-212 (1978). Most of such compounds are structural variations of 1-amino-3-aryloxy-2-propanol.


5 Heretofore, the emphasis in β -blocker research has been to develop compounds which can be administered to cardiac patients over long periods of time. However, often it is desirable in the critical care setting to quickly reduce heart work or improve rhythmicity during a cardiac crisis, e.g., during or shortly after a myocardial infarction. Conventional β -blocking agents can be employed for such treatment, but their duration of action may be much longer than desired by the physician. A β -blocking agent possessing a long duration of action does not allow precise control of heart work or prompt reversal of the β -blocking effect, which may be required in a critical care setting. For instance, if heart output becomes dangerously low, it is desirable to quickly reduce or eliminate β -blocking activity. The lingering activity of available β -blocking agents can be counter-productive and can greatly complicate the therapeutic decisions required of the physician during such critical care of cardiac patients.

Summary of the Invention

In accordance with the present invention, disclosed herein is a method for the treatment or prophylaxis of cardiac disorders in a mammal comprising administering to such mammal a short-acting compound of the formula:




30 wherein Y is a straight or branched carbon chain of from 1 to about 10 carbon atoms or aralkyl of from 8 to about 20 carbon atoms; R is lower alkyl, lower alkenyl, lower alkynyl, aryl or aralkyl; x is an integer from 1 to about 3; Ar is unsubstituted aromatic or aromatic substituted with lower alkyl, lower alkenyl, lower alkynyl, lower alkoxy, halogen, acetamido, amino, nitro, lower alkylamino, hydroxy, lower hydroxyalkyl, cyano, or a group of the formula



wherein n is an integer from 0 to about 10; or a pharmaceutically acceptable salt thereof.

Detailed Description of the Invention

5 In accordance with the present invention, it has been discovered that compounds having an ester function in external amine substituents possess β -adrenergic blocking activity and have a short duration of action. Such compounds may also contain more than one ester group in the same molecule. the compounds of the present invention are
10 represented by the formula:




15 Y may be a straight or branched carbon chain of from 1 to about 10 carbon atoms, e.g., methylene, ethylene, propylene, 2-ethylhexylene, 1,1-dimethylethylene, and the like or aralkyl of, from 8 to about 20 carbon atoms, such as dialkylene phenyl, e.g., 4-ethylenebenzyl,
20 1-propylene-(4-naphthyl)-2-n-butyl, and the like.

R may be lower alkyl of from 1 to about 10 carbon atoms, such as methyl, propyl, t-butyl, 3-propylheptyl, and the like; lower alkenyl of from 2 to about 10 carbon atoms, such as ethenyl, propenyl, 4-ethyl-2-
25 hexenyl, and the like, lower alkynyl of from 2 to about 10 carbon atoms, such as ethynyl, propynyl, 4-ethyl-3-octynyl, and the like; aryl of from 6 to about 10 carbon atoms such as phenyl, 2-tolyl, 2-methoxyphenyl, naphthyl, and the like or aralkyl, wherein the alkyl portion contains from 1 to about 10 carbon atoms and the aryl portion contains
30 from 6 to about 10 carbon atoms, such as benzyl, phenethyl, 1-naphthylpropyl, 3,4-dimethoxyphenethyl, and the like.

The amine substituent may contain one or more ester groups, thus x
35 is an integer from 1 to about 3 provided that when x is greater than 1, different occurrences of the-COOR group may be the same or different.

Ar represents substituted or unsubstituted aromatic, including monocyclic, polycyclic, and heterocyclic ring systems. Aromatic substituents include lower aryl, of from 1 to about 10 carbon atoms, lower alkenyl, of from 2 to about 10 carbon atoms, lower alkynyl, of from 2 to about 10 carbon atoms, lower alkoxy of from 1 to about 10 carbon atoms, halogen, acetamido, amino, nitro, lower alkylamino, of from 1 to about 10 carbon atoms, hydroxy, lower hydroxyalkyl of from 1 to about 10 carbon atoms, cyano, or a group of the formula

10



wherein n is an integer from 0 to about 10. When two or more groups of the same designation occur in the same formula, those groups are not necessarily identical. The compounds described herein are not limited to any particular stereoisomeric configuration.

In preferred compounds, Y is a straight or branched carbon chain of from 1 to about 6 carbon atoms or aralkyl of from 8 to about 12 carbon atoms. Most preferably, Y is a straight or branched carbon chain of from 1 to about 4 carbon atoms. R is preferably lower alkyl of from 1 to about 5 carbon atoms, lower alkenyl of from 2 to about 5 carbon atoms, lower alkynyl of from 2 to about 5 carbon atoms, aryl of from 6 to about 8 carbon atoms, or aralkyl, wherein the alkyl portion contains from 1 to about 5 carbon atoms and the aryl portion contains from 6 to about 10 carbon atoms. Most preferably, R is lower alkyl of from 1 to about 4 carbon atoms or aralkyl, wherein the alkyl portion contains from 1 to about 4 carbon atoms and the aryl portion contains from 6 to about 8 carbon atoms. Particularly preferred R groups are methyl and ethyl. The integer x is preferably 1 or 2; most preferably 1.

Ar is preferably unsubstituted aromatic or aromatic substituted with lower alkyl of from 1 to about 5 carbon atoms, lower alkenyl of from 2 to about 5 carbon atoms, lower alkynyl of from 2 to about 5 carbon atoms, lower alkoxy of from 1 to about 5 carbon atoms, fluoro, chloro, acetamido, amino, nitro, lower alkylamino of from 1 to about 5

carbon atoms, hydroxy, lower hydroxyalkyl of from 1 to about 5 carbon atoms, cyano, or a group of the formula




wherein n is an integer from 0 to about 5. Ar is more preferably unsubstituted phenyl or phenyl substituted with lower alkyl of from 1 to about 5 carbon atoms, fluoro, chloro, nitro or a group of the formula



15 wherein n is an integer of from 1 to about 5 and R is lower alkyl of from 1 to about 5 carbon atoms. Most preferably, Ar is 2-alkylphenyl, eg. 2-methylphenyl.


The compounds of this invention may be administered as their pharmaceutically acceptable acid addition salts, e.g., as the hydrochloride, sulfate, phosphate, gluconate, tartrate, etc.

The compounds of the present invention may be prepared by a number of reaction procedures. The following four reaction schemes have been employed. In all of the reaction schemes, an aryl ether epoxide is used as the starting material. The aryl ether epoxide is prepared from an appropriately derivatized aryl hydroxy compound as follows:




The aryl ether epoxide may then be reacted in the following manner to provide the desired product:


Scheme I




Scheme II



Scheme III



Scheme IV



The compounds of this invention are advantageously administered parenterally, e.g., by intravenous injection or intravenous infusion. Formulations for intravenous injection preferably include the active compound as a soluble acid addition salt in a properly buffered isotonic solution.

The dosage administered to a patient and the duration of infusion will depend upon the patient's needs and the particular compounds employed. For short periods of infusion, e.g., less than about three hours, the duration of effect is thought to be determined by both metabolic effects and distribution phenomena. For relatively long periods of infusion, e.g., greater than about three hours, the duration of effect is thought to depend largely on metabolic effects. Accordingly, although the present methods and compounds are generally useful for short term infusion therapy, certain compounds are preferred for longer durations of infusion. This principle is demonstrated by referenced to the 40 minute and three hour infusion studies described in Examples LX-LXXIV. The compounds have been found to be generally non-toxic within conventional dosage ranges. Dosages of from about 0.001 to about 100 mg. per kg. of body weight per hour are generally employed,


with preferred dosages ranging from about 0.01 to about 10 mg. per kg. of body weight per hour.

The compounds of the present invention have a relatively short duration of action compared to conventional β -blockers. In vitro studies in human whole blood indicate that the ester functions are subject to enzymatic cleavage. Compounds of the present invention in which the aromatic portion, Ar, is also substituted with an ester-containing group, have two or more potentially labile sites for enzymatic hydrolysis. Thus β -blocking activity can be carefully controlled by regulating dosage size and rate of administration. The time required for substantially complete disappearance of the β -blocking effects of the compounds of the present invention ranges from about 5-10 minutes to about 1 hour or more. Generally, it is preferred that the recovery is accomplished within about 10-15 minutes. A short acting β -blocker can advantageously be infused at a rate sufficient to provide the desired action, e.g., titrated to the specific patient's needs, and such action can be promptly discontinued by stopping the infusion. Thus, the method of the present invention provides a very useful therapeutic alternative in the treatment or prophylaxis of cardiac disorders.

The present invention is further illustrated by the following examples which are not intended to be limiting.

Example I

This example describes procedures producing the following compound:



2,3-Epoxypropoxybenzene

A mixture of 9.4 gm (0.1 mole) of phenol, .28 gm (0.2 mole) of potassium carbonate and 30 mL (0.4 mole) of epichlorohydrin in 250 mL acetone was heated to reflux for 12 hours. The reaction medium was then


filtered and evaporated leaving an oil which was taken up in toluene and successively washed with 100 mL water, 2 x 100 mL 1.0 N sodium hydroxide and 2 x 100 mL water. The toluene phase was dried with magnesium sulfate and evaporated to provide a clear oil which was chromatographed
 5 on a Prep-500 silica gel column employing hexane: ethyl acetate (9:1) as the mobile phase. Collection of the major peak and evaporation of solvent provided 9 gm (60%) of a clear oil whose NMR and IR spectra were consistent with the assigned structure.

10 Ethyl N-(2-Hydroxy-3-phenoxypropyl) glycinate Oxalate

A solution of 20 gm (0.14 mole) of glycine ethyl ester hydrochloride and 40 gm K_2CO_3 in 100 mL of water was extracted with ether (5 x 100 mL). The ethereal phase was then dried with Na_2SO_4 and evaporated under reduced pressure at a temperature not exceeding 40°C
 15 to provide 10.5 gm (71%) of glycine ethyl ester free amine. The glycine ethyl ester free amine (0.10 mole) was used immediately by reacting with 4.0 gm (0.03 mole) of 2,3-epoxypropoxybenzene in refluxing ethanol (50 mL). After 4 hours the reaction medium was evaporated under reduced pressure and the resulting oil taken up in 50 mL toluene and washed with
 20 4 x 50 mL water. The organic phase was dried with $MgSO_4$ and evaporated to a yellow oil. An analytical sample of the free amine was obtained by crystallization from ethyl acetate: mp 49-50°. The elemental analysis of this product was consistent with the formula, $C_{13}H_{19}NO_4$. The major portion of this oil was converted to its oxalate salt and crystallized
 25 from ethanol-ether to provide 0.8 gm (8%): mp 144-145°C. The NMR and IR spectra and elemental analyses conformed to the assigned structure.

Example II

This example describes the procedures for producing a compound of
 30 the formula



1-Naphthyl-2,3-epoxypropyl Ether


The experiment of Example I for preparing 2,3-epoxypropoxybenzene was repeated in all essential details, except that 1-naphthol was substituted for phenol, and the final product was isolated by vacuum
 5 distillation: bp 112-114°C (p=0.25 mm Hg). The NMR and IR spectra and elemental analyses conformed to the assigned structure.

Ethyl N-[[2-Hydroxy-3-(1-naphthoxy)propyl]glycinate Oxalate Hemihydrate

A mixture of 4.0 gm (0.02 mole) of 1-(2,3-epoxypropoxy)-
 10 naphthalene, 5.6 gm (0.04 mole) of glycine ethyl ester hydrochloride and 5.5 mL (0.04 mole) triethylamine in 50 mL of ethanol was heated to reflux for 2 hours. The reaction medium was then evaporated under reduced pressure and the resulting oil taken up in 50 mL toluene and washed with 2 x 50 mL water. The organic phase was dried over MgSO₄ and
 15 evaporated under reduced pressure. The resulting oil was crystallized as its oxalate salt from water and provided 1.1 gm (14%) of product: mp 161-162°C. The NMR spectrum was consistent with the assigned structure, and the elemental analysis was consistent with the formula C₁₉H₂₃NO₈.1/2H₂O. An anhydrous analytical sample was also obtained by
 20 crystallization from acetone: mp 169-170°C. The NMR spectrum of the analytical sample was also consistent with the assigned structure, and the elemental analysis was consistent with the formula C₁₉H₂₃NO₈.

Example III

25 This example describes procedures for producing the following compound:

1-Succinimido-3-phenoxy-2-propanol

A mixture of 15 gm (0.1 mole) of 2,3-epoxypropoxybenzene (prepared as described in Example I) and 9.9 gm (0.1 mole) of
 35 succinimide in 100 mL ethanol having 10 drops of pyridine was heated to reflux for 4 hours. After standing 24 hours at room temperature a white crystalline product separated. This material was collected, air-dried

0053434

and recrystallized from 700 mL ethyl acetate:hexane (6:1) to give 18 gm (72%) of white crystals: mp 130°. The NMR spectrum and the elemental analyses were consistent with the assigned structure.

5 1-Amino-3-phenoxy-2-propanol Hydrochloride

1-Succinimido-3-phenoxy-2-propanol (16 gm, 0.06 mole) was dissolved in 100 mL conc. HCl and 100 mL ethanol and heated to reflux for 6 hours. After the reaction, the mixture was evaporated to a white residue which was then taken up in 25 mL water and washed with 3 x 50 mL
10 ether. The aqueous phase was then evaporated and the white-residue recrystallized from ethanol to provide 8.3 gm (69%) of white crystals: mp 226-228°. The NMR spectrum was consistent with the assigned structure and the elemental analysis was consistent with the molecular formula $C_9H_{14}NO_2Cl$.

15


Ethyl 3-[N-[2-Hydroxy-3-phenoxy)propyl] amino]propionate Oxalate

A mixture of 3.4 gm (0.02 mole) of 1-amino-3-phenoxy-2-propanol hydrochloride, 2.6 mL (0.02 mole) of ethyl 3-bromopropionate and 2.8 mL (0.02 mole) of triethylamine in 20 mL of ethanol was heated to reflux
20 for 12 hours. The reaction medium was then filtered and evaporated and the residue taken up in 25 mL water/50 mL ether. The phases were separated and the ether phase washed twice with 25 mL water. The ethereal phase was then dried with magnesium sulfate and evaporated to provide 1.86 gm of a clear oil. A 1.65 gm (0.0062 mole) portion of this
25 oil in 5 mL ethanol was then added to 0.78 gm (0.0062 mole) of oxalic acid dihydrate in 15 mL ethanol and after standing at room temperature the oxalate salt was produced as a crystalline solid. The material was recrystallized from acetone to provide 1.1 gm (16%) white crystals: mp 137-138°. The NMR spectrum was consistent with the assigned structure
30 and the elemental analysis was consistent with the molecular formula $C_{16}H_{23}NO_8$.

Example IV

This example describes a procedure for preparing a compound of
35 the formula

0053434




5 Ethyl 4-[N-[(2-Hydroxy-3-phenoxy)propyl]aminobutyrate

A mixture of 10 gm (0.10 mole) of 4-aminobutyric acid, 6.6 mL of 2,3-epoxypropoxybenzene (0.05 mole) (prepared as described in Example I) and 4.0 gm (0.10 mole) of NaOH in 160 mL aqueous dioxane (1:3) was heated to reflux for 4 hours. After cooling, 100 mL of water was added and the aqueous medium washed with 400 mL ether. The aqueous phase was acidified to pH 1 by adding concentrated HCl and then evaporated to a semi-solid residue which was extracted with ethyl acetate. This process removed 5.3 gm (95%) of NaCl side product. Evaporation of ethyl acetate provided the crude amino acid product as an oil which was immediately esterified with 500 mL ethanol utilizing a Soxhlet Extractor charged with 250 gm of activated 3A molecular sieves and employing a 96-hour reaction time. Concentration of the ethanol and treatment with ether provide a crystalline material which was subsequently recrystallized from ethyl acetate to provide 4.1 gm (25%) of product: mp 109-100°C. The NMR spectrum was consistent with the assigned structure, and the elemental analysis was consistent with the formula $C_{15}H_{24}NO_4Cl$.

Example V

25 This example describes the experimental procedures for producing the following compound:



35 Ethyl N[(2-Hydroxy-3-phenoxy)propyl]tyrosinate

A mixture of 1.4 mL (0.01 mole) of 2,3-epoxypropoxybenzene (prepared as described in Example I) and 2.1 gm (0.01 mole) of tyrosine

in 10 mL ethanol was heated to reflux for 4 hours. After the reaction, the mixture was evaporated to a thick clear oil which was then dissolved in 50 mL toluene and partitioned with 2 x 40 mL water. The organic phase was then dried with magnesium sulfate and evaporated to an oil.


5 This oil was taken up in 15 mL of ethanolic HCl and treated with 175 mL ether. An oil was gradually produced from this solution after cooling. This oil was taken up in ethyl acetate and upon evaporation of this solvent, and amorphous solid was obtained: 0.5 gm (14%); mp 60-70°. The NMR spectrum was consistent with the assigned structure and the

10 elemental analysis was consistent with the molecular formula $C_{20}H_{26}NO_5Cl$.

Example VI

This example describes a procedure for preparing a compound of

15 the formula



Ethyl 3[N-[3-[4-chlorophenoxy]-2-hydroxy]propyl]amino] propionate Hydrochloride

A solution of 9.2 gm (50 mmol) of 4-chloro-1-(2,3-epoxypropoxy) benzene and 5.5 mL (50 mmol) of benzylamine in 125 mL of ethanol was

25 heated to reflux for 4 hours (a 10 mL aliquot was then treated with concentrated HCl and ether to provide an analytical sample of the intermediate secondary benzylamine hydrochloride as a white crystalline product: mp 169-170°C). After cooling the reaction mixture, 6 mL (47


30 mmol) of ethyl 3-bromopropionate and 6.5 mL (47 mmol) of triethylamine were added and the mixture heated to reflux for another 10 hours. The reaction medium was then evaporated and the residue taken up in 50 mL toluene - 50 mL water. The organic phase was washed an additional two

35 times with 50 mL portions of water and then dried over $MgSO_4$ and evaporated to provide the Ethyl 3-[N-benzyl-N[3-[(4-chlorophenoxy)-2-hydroxy]propyl]amino]propionate intermediate as an oil which was characterized by NMR spectroscopy. This oil was used directly in the

next reaction by redissolving it in 100 mL of ethanol, adding 7 mL (100 mmol) of acetyl chloride, 100 mg of 10% Pd-C and hydrogenating under 50 psi for 20 minutes. The reaction medium was then filtered and evaporated under reduced pressure to provide the product as an oil which yielded 6.5 gm (41% overall yield) of white crystals from ethanol-ether: mp 119-120°C. The NMR spectrum and elemental analysis were consistent with the assigned structure.

Example VII

This example described procedures for preparing a compound of the formula



Methyl 3(4-Hydroxyphenyl)propionate

A solution of 300 gm (1.81 mole) of 3-(4-hydroxyphenyl) propionic acid in 1 liter of anhydrous methanol containing 10 drops of concentrated H_2SO_4 was heated to reflux for 72 hours in a Soxhlet Extractor charged with 200 gm of 3A molecular sieves (Linde 3A, 1/16 pellets). The reaction medium was then evaporated under reduced pressure and the resulting oil taken up in 750 mL of toluene and washed with three 500 mL portions of water. The toluene phase was then dried with MgSO_4 and evaporated under reduced pressure to provide 228.4 gm (70%) of a clear oil which was characterized by NMR spectroscopy and utilized directly in the next step without additional purification.

Methyl 3[4-(2,3-Epoxypropoxy)phenyl]propionate

A mixture of 228 gm (1.27 mole) of methyl 3-(4-hydroxyphenyl)propionate, 263 gm (1.90 mole) of K_2CO_3 and 298 mL (3.80 mole) of epichlorohydrin in 2 liters of acetone was stirred and heated to reflux for 20 hours. The reaction medium was then filtered and evaporated under reduced pressure. The resulting oil was taken up in 1 liter of

0053434


toluene and washed consecutively with 500 mL water, 2 x 500 mL 1N NaOH and 2 x 500 mL water. The toluene phase was then dried over MgSO_4 and evaporated under reduced pressure to provide a clear oil which was further purified by vacuum distillation. The final yield of purified oil was 131.2 gm (44%): bp 156° (p=0.4 mm Hg). The NMR and IR spectra of the product were consistent with the assigned structure and the elemental analysis was consistent with the formula $\text{C}_{13}\text{H}_{16}\text{O}_4$.

10 Ethyl 3-[N-[2-Hydroxy-3-[4-[2-(methoxycarbonyl)ethyl]phenoxy]propyl]-amino]propionate Hydrochloride

A mixture of 5 gm (0.02 mole of methyl 3-[4-(2,3-epoxypropoxy)-phenyl]propionate, 3 gm (0.02 mole) of ethyl 3-aminopropionate hydrochloride and 2.8 ml (0.02 mole) of triethylamine in 25 mL of isopropanol was heated to reflux for 4 hours. The reaction medium was then cooled and the triethylamine hydrochloride side-product which crystallized was removed by filtration. The mother liquor was then evaporated under reduced pressure and the resulting residue taken up in ethanol and treated with ethereal HCl to provide 1 gm (12%) as a crystalline solid: mp $110-111^\circ$. The NMR spectrum was consistent with the assigned structure and the elemental analysis was consistent with the formula $\text{C}_{18}\text{H}_{28}\text{NO}_6\text{Cl}$.

Example VIII

25 This example describes procedures for the preparation of a compound of the formula



Methyl 2-(2,3-Epoxypropoxy)benzoate

35 The procedure of Example VII for producing methyl 3-[4-(2,3-epoxypropoxy)phenyl]propionate was repeated in all essential details, except methyl salicylate was substituted for methyl 3-(4-hydroxyphenyl)propionate. The boiling point of the product was 148° (p=75 μ).

The NMR spectrum was consistent with the assigned structure and the elemental analysis with the formula $C_{11}H_{12}O_4$.

Ethyl 3-[N-[2-Hydroxy-3-[2-(methoxycarbonyl)phenoxy]propyl]amino] -
5 propionate Oxalate Hemihydrate

The procedure for Example VII for producing ethyl 3-[N-[2-hydroxy-3-[4-[2-(methoxycarbonyl)ethyl]phenoxy]propyl]amino]propionate hydrochloride was repeated in all essential details except methyl 2-(2,3-epoxypropoxy) benzoate was substituted for 3[4-(2,3-epoxypropoxy)phenyl]propionate and the product was crystallized as its
10 oxalate salt from 2-propanol ether and then recrystallized from acetone. Approximately 2 gm (25%) of product was obtained having a melting point of 90-91°C. The NMR spectrum was consistent with the assigned structure and the elemental analysis was consistent with the formula
15 $C_{18}H_{25}NO_{10} \cdot 1/2H_2O$.

Example IX


This example describes the preparation of 1-(2,3-epoxypropoxy)-2-methylbenzene, which may be used as a starting material for certain
20 compounds described herein. The procedure is representative and may be modified to provide starting material for a variety of compounds. A mixture of 52 mL (0.5 mole) of ortho-cresol, 103 gm (0.75 mole) of K_2CO_3 and 117 mL (1.5 mole) of epichlorohydrin in 600 mL of acetone was heated to reflux for 16 hours. The reaction medium was then filtered and
25 evaporated under reduced pressure. The resulting oil was taken up in 400 mL of toluene and washed consecutively with 200 mL of water, 2 x 200 mL of 1.0 N aq. sodium hydroxide and 200 mL of water. The organic phase was then dried over $MgSO_4$ and evaporated under reduced pressure. The resulting oil, 54 gm (65%), was utilized directly in the next step
30 without additional purification. The product was characterized by NMR spectroscopy, and the spectrum was consistent with the assigned structure.

Examples X - XX

35 These examples describe the preparation of compounds identified in Table I. The compounds were prepared utilizing the procedure of

Example VI in all essential details, except 1-(2,3-epoxypropoxy)-2-methylbenzene was substituted for 4-chloro-1-(2,3-epoxypropoxy)benzene and the appropriate bromocarboxylic acid ester was substituted for ethyl 3-bromopropionate. Each of the compounds was characterized by NMR spectroscopy and elemental analysis.

Table I




15







Example	Y	R	Salt	Melting Point	Yield (%)
X	-CH ₂ -CH ₂ -	-CH ₂ CH ₃	HCl	82-84°C	44
XI	-(CH ₂) ₃ -	-CH ₂ CH ₃	HCl	77-78°C	16
20 XII	-(CH ₂) ₄ -	-CH ₂ CH ₃	HCl	102-103°C	46
XIII		-CH ₂ CH ₃	Free Amine	102-103°C	20
XIV		-CH ₂ CH ₃	Oxalate	89-92°C	18
XV	-CH ₂ -	-CH ₂ CH ₃	HCl	144-145°C	15
XVI	-CH ₂ -	-CH ₃	HCl	179-180°C	12
25 XVII	-CH ₂ -CH ₂ -	-CH ₃	Oxalate	152-153°C	10
XVIII	-CH ₂ -CH ₂ -		HCl	148-149°C	16
XIX	-CH ₂ -CH ₂ -	F	HCl	142-144°C	14
30 XX	-CH ₂ -CH ₂ -	-CH ₂ CF ₃	Hemi-Oxalate	133-134°C	5

Examples XXI - XXVI

These examples describe the preparation of the compounds identified in Table II. The compounds were prepared utilizing the procedure of Example VI in all essential details, except the appropriate 2,3-epoxypropoxyaryl compound was substituted for 4-chloro-(2,3-epoxypropoxy)benzene. Each of the compounds was characterized by NMR spectroscopy and elemental analysis.

Table II




Example	Ar	Salt	Melting Point	Yield %
XXI		Oxalate	134-137°C	15
XXII		HCl	94-95°C	21
XXIII		Oxalate	127-128°C	18
XXIV		Oxalate	129-132°C	22
XXV		HCl	117-119°C	20
XXVI		Hemi-Oxalate	148-149°C	5

Example XXVII - XXVIII

These examples describe the preparation of the compounds identified in Table III. The compounds were prepared utilizing the procedure of Example III in all essential details, except the appropriate bromocarboxylic acid ester was substituted for ethyl 3-bromopropionate, and 1-(2,3-epoxypropoxy)2-methylbenzene was substituted for 2,3-epoxypropoxybenzene. The compounds were identified by NMR spectroscopy and elemental analysis.

Table III




10

Example	Y	R	Melting Point	Salt	Yield (%)
XXVII	-CH ₂ -	-CH ₂ CH ₃	137-138°C	Oxalate	14
XXVIII	-CH ₂ CH ₂ -	<p>15</p>	125-129°C	Hemi-Oxalate	23

Example XXIX

The procedure of Example IV was repeated in all essential details to produce a compound of the formula

20



25 except 1-(2,3-epoxypropoxy)naphthalene was substituted for 2,3-epoxypropoxybenzene, 3-aminopropionic acid was substituted for 4-aminobutyric acid, and methanol was substituted for ethanol in the esterification step. The product was crystallized in 10% yield as its oxalate salt and had a melting point of 180°C. The NMR spectrum and elemental analysis were consistent with the assigned structure.

30


Example XXX

This example describes procedures for the preparation of a compound of the formula

35

0053434

5



Diethyl 2-Amino-[N-[2-Hydroxy-3-(2-methylphenoxy)propyl]]propanedioate Hydrochloride


A mixture of 10g (0.061 mole) of 2-methyl-1-(2,3-epoxypropoxy)-
 10 benzene, 12.9 gm (0.061 mole) of diethylaminomalonate hydrochloride and
 6.2 gm (0.061 mole) of triethylamine in 100 mL of ethanol was heated to
 reflux for 24 hours. The reaction medium was then evaporated under
 reduced pressure and the residue treated with ether. The solid tri-
 ethylamine hydrochloride side product was then removed by filtration.
 15 The mother liquor was then treated with HCl gas and provided 2.7 (12%)
 of white crystals: mp 105-6°. The NMR spectrum was consistent with the
 assigned structure, and the elemental analysis was consistent with the
 formula $C_{17}H_{26}NC_2O_6$.

20

Example XXXI

This example describes procedures for the preparation of a com-
 pound of the formula:

25



30

Diethyl 2-Amino-[N-[2-hydroxy-3-(4-(2-carbomethoxyethyl)phenoxy)propyl]]propanedioate Hydrochloride

A mixture of 11.8 gm (0.05 mole) of methyl 3-[4-(2,3-epoxy-
 propoxy)phenyl]propionate, 10.6 gm (0.05 mole) of diethyl
 2-aminomalonate hydrochloride and 7 mL (0.05 mole) of diethyl 2-
 35 aminomalonate hydrochloride and 7 mL (0.05 mole) of triethylamine in
 100 mL of isopropanol was heated to reflux for 4 hours. The reaction

medium was then filtered and evaporated under vacuum. The resulting oil was taken up in 100 mL toluene and washed with 3 x 50 mL water. The organic phase was then dried over MgSO_4 and evaporated under vacuum. The free amine oil was taken up in ethanol and converted to its hydrochloride salt by adding ethereal HCl . Approximately 4 gm (20%) of white crystals were obtained: mp 144-145°. The NMR spectrum was consistent with the assigned structure and the elemental analysis was consistent with the formula $\text{C}_{20}\text{H}_{30}\text{NO}_8\text{Cl}$.

10

Examples XXXII - LIX

Several of the compounds of the present invention were tested for β -blocking activity in vitro using guinea pig right atria and guinea pig tracheal strips mounted in a tissue bath containing oxygenated (95% O_2 -5% CO_2) Krebs physiological salt solution at 37°C. Each tissue was suspended between a fixed glass rod and a Statham Universal Transducer connected to a Beckman recorder. Atria were allowed to beat spontaneously under a loading tension of approximately 0.5 gm. Intrinsic depressant or stimulant activity was determined for each compound by progressively increasing concentrations in the tissue baths at 60-minute intervals. Tissues were not washed between increments. The maximum concentration showing little or no cardiodepressant activity was chosen for blockade experiments. Changes in rate in response to isoproterenol were measured in the absence and presence of test compounds. Spiral strips of guinea pig trachea were suspended under 5 gm resting tension and incubated with phentolamine, tropolone and cocaine. Active tension was generated by addition of carbachol ($3.0 \times 10^{-7}\text{M}$) and decreases in tension in response to isoproterenol were quantitated. Cumulative concentration-response curves were produced with isoproterenol both before and after 60 minute incubation of test compounds with atria and trachea. The blocking potency of test compounds was estimated by computing pA_2 values ($-\log K_B$) by the method of Furchgott, The Pharmacological Differentiation of Adrenergic Receptors, Ann. N.Y. Acad. Sci.; 139: 553-570 (1967). Comparison of blockade of right atrial and tracheal responses to isoproterenol permitted assessment of cardioselectivity of test compounds; i.e., cardioselective compounds are relatively more effective in blocking atrial rate than tracheal force responses to isoproterenol. The degree

35

0053434

of cardioselectivity was estimated from the ratio, K_B tracheal/ K_B atrial ($10^{(PA_{2\text{atria}} - PA_{2\text{trachea}})}$). A ratio greater than one indicates cardioselectivity. Test drugs were dissolved in distilled water and added to the bath (30 mL) in a volume of 10 or 100 μ L. The results of the in vitro tests are contained in Table IV. All of the test compounds were active β -blockers.

Examples LV - LXXIV

The duration of β -blockade was determined in vivo using pentobarbital-anesthetized dogs instrumented for measurement of heart rate using a Beckman cardi tachometer triggered electronically by a phasic aortic blood pressure signal. Both vagus nerves were severed in the cervical region and the animals were mechanically ventilated. Two experimental designs were used. The first employed a 40-minute infusion of test compound and the second used a 3-hour infusion of test compound. In the 40-minute model, isoproterenol was infused into a foreleg vein at the rate of 0.5 μ g/kg/min to induce a β -receptor mediated tachycardia. Various doses of test compound are then infused into a femoral vein over a period of 40 minutes. This infusion was then terminated and recovery from blockade was quantitated. The percent inhibition of the heart rate response to isoproterenol after 40 minutes of infusion of the test compound was computed along with the total cumulative doses received of the 40-minute period. This cumulative dose is expressed as mg/kg and is an indication of potency. The time period required for 80% recovery of heart rate response for each dose of test drug was also measured to quantitate duration of action. To facilitate comparison of data between animals, the data for potency and duration of action were normalized to a level of 50% inhibition of the isoproterenol response via least squares regression of data from each animal. Test compounds were dissolved in 0.9% NaCl and infused at a rate of 0.05 mL/kg/min or less. In the 3-hours infusion model, bolus doses of isoproterenol (0.4 μ g/kg) were used to assess the degree of β -blockade and recovery from β -blockage after termination of the infusion. The doses were spaced at 10-minute intervals and were given before, during and following the infusion of test compounds. The infusion rate was adjusted so that at the end of the 3-hour infusion period the degree of isoproterenol inhibition averaged about 50% of

control. The results of isoproterenol inhibition averaged about 50% of control. The results of the 40-minute infusion are shown in Table V, and the results of the 3-hour infusion are shown in Table VI.

5

Example LXXV - LXXXII

10 These examples describe experiments which demonstrate the disappearance of the compounds of the present invention in vitro in human whole blood, dog whole blood, and dog liver homogenate. The rate of disappearance of a compound is expressed as the half-life ($T_{1/2}$), which is the time period in which one-half of the initial amount of compound tested disappears. In each experiment, 1 mL of a solution containing 50 ug of the test compound was added to 1 mL of whole blood or 1 mL of a 33% (w/v) liver homogenate. The samples were incubated in a Dubnoff shaking metabolic incubator for 2.5, 5.0, 10.0, 20.0, 30.0 and 15 60.0 minutes at 37°C. At the designated time periods, the test mixtures were removed from the incubator and transferred to a 0°C ice bath. Acetonitrile (2 mL) was immediately added and the mixtures were mixed to stop enzymatic hydrolysis. Zero time samples were prepared by adding 2 mL of acetonitrile to denature the proteins prior to addition of the 20 test compounds. After centrifugation to sediment denature proteins, 2 mL of the supernatant was removed and analyzed by high pressure liquid chromatography, using a mobile phase of 60% acetonitrile/40% 0.5M sodium phosphate buffer (pH 6.6), a U.V. detector and Waters μ Bondapak Phenyl column. The half life of each test compound was determined 25 graphically by plotting the decrease in concentration as a function of time. The results of the experiments are shown in Table VII.

0053434

Table IV

Example	Test Compound (Numerical Designation Indicates Previous Example Which Describes Preparation of Compound)	pA ₂		Cardioselectivity
		Atria	Trachea	
XXXII	I	5.7	5.6	1
XXXIII	II	6.2	6.2	1
XXXIV	III	7.3	5.9	25
XXXV	IV	6.6	6.0	4
XXXVI	V	inactive	inactive	-
XXXVII	VI	6.4	5.6	6
XXXVIII	VII	5.9	<	8
XXXIX	VIII	6.6	6.6	>
XL	X	7.5	7.4	1.0
XLI	XI	7.4	6.7	1
XLII	XII	7.8	7.4	5
XLIII	XIII	5.1	-	2.5
XLIV	XIV	5.2	-	-
XLV	XV	6.6	5.9	-
XLVI	XVI	6.9	6.7	5
XLVII	XVII	7.5	7.0	1.6
XLVIII	XVIII	6.4	-	1.6
XLIX	XIX	5.7	-	-
L	XX	5.7	-	-
LI	XXI	7.4	6.6	-
LII	XXII	7.4	6.7	6
LIII	XXIII	6.7	6.5	5
LIV	XXIV	6.4	6.7	1.6
LV	XXV	5.8	4.9	.5
LVI	XXVI	6.1	-	8
LVII	XXVII	5.9	6.2	-
LVIII	XXVIII	7.0	6.6	0.5
LIX	XXIX	7.1	7.6	2.5
Propranolol		8.7	8.9	0.3
Practolol		6.6	5.8	0.6
				6.0

0053434

Table V

Example	Test Compound (Numerical Designation Indicates Previous Example Which Describes Preparation of Compound)	Potency (mg/kg/40 min)	80% Recovery Time (Minutes)	No. of Experiments
LX	I	very low	-	4
LXI	II	-	53+11	-
LXII	III	12.6+4.1	29+13	4
LXIII	IV	16.9+9.1	17+4	3
LXIV	XXV	17.2+1.2	11+2	4
LXV	X	0.99+0.27	14+2	5
LXVI	XI	3.1+0.7	15+2	4
LXVII	XII	6.8+3.7	11+2	3
Propranolol			46	
Practolol			61	

0053434

Table VI

Example	Test Compound (Numerical Designation Indicates Previous Example Which Describes Preparation of Compound)	Potency (mg/kg/180 min)	%I*	Recovery Time (Minutes)	No. of Experiments
LXVIII	XVII	4.4	56+2	27+4	4
LXIX	VIII	27.4+6.0	53+2	10+2	5
LXX	XXVIII	167.7	85	>60	1
LXXI	X	6.8	69+5	>60	3
LXXII	XXII	8.5	77	>60	3
LXXIII	XXI	10.9	77	>60	2
LXXIV	XVIII	54+14	54+8	>60	3

No appreciable
recovery after
several hours

No appreciable
recovery after
several hours

Propranolol

Practolol

*Percent inhibition of heart rate response to isoproterenol

0053434


Table VII

Example	Test Compound*	$T_{1/2}$ (Minutes)		
		Human Whole Blood	Dog Whole Blood	Dog Liver Homogenate
LXXV	XII	93+30	152+71	8.8+6
LXXVI	XI	117+39	>180	-
LXXVII	XXII	118+26	40+15	3+3
LXXVIII	XXI	127+50	74+21	4+3
LXXIX	XVII	148+2	52+9	9+8
LXXX	X	155+9	180+0	3+1
LXXXI	XXV	158+8	>180	4+2.5
LXXXII	XV	>180	>180	9+6


*Numerical Designation Indicates Previous
Example which Describes Preparation of Compound

CLAIMS

1. A compound of the formula



wherein Y is a straight or branched carbon chain of from 1 to about 10
 10 carbon atoms or aralkyl of from 8 to about 20 carbon atoms; R is lower
 alkyl, lower alkenyl, lower alkynyl, aryl, or aralkyl; x is an integer
 from 1 to about 3, provided that when x is greater than 1, different
 occurrences of the -COOR group may be the same or different; Ar is
 15 unsubstituted aromatic or aromatic substituted with lower alkyl, lower
 alkenyl, loweralkynyl, lower alkoxy, halogen, acetamido, amino, nitro,
 lower alkylamino, hydroxy, lower hydroxyalkyl, cyano, or a group of the
 formula




wherein n is an integer from 0 to about 10; or a pharmaceutical
 acceptable salt thereof.

- 25 2. The compound of Claim 1 wherein R is lower alkyl of from 1
 to about 10 carbon atoms, lower alkenyl of from 2 to about 10 carbon
 atoms, lower alkynyl of from 2 about 10 carbon atoms, aryl of from 6 to
 about 10 carbon atoms or aralkyl, wherein the alkyl protion contains
 from 1 to about 10 carbon atoms and the aryl portion contains from 6 to
 30 about 10 carbon atoms.

3. The compound of claim 1, wherien Y is a straight or
 branched carbon chain of from 1 to about 6 carbon atoms or aralkyl of
 from 8 to about 12 carbon atoms and R is lower alkyl of from 1 to about 5
 35 carbon atoms, lower alkenyl of from 2 to about 5 carbon atoms, lower
 alkynyl of from 2 to about 5 carbon atoms, aryl of from 6 to about 10

carbon atoms or aralkyl wherein the alkyl portion contains from 1 to about 5 carbon atoms and the aryl portion contains from 6 to about 10 carbon atoms.

- 5 4. The compound of claim 3 wherein Ar is unsubstituted aromatic or aromatic substituted with lower alkyl of from 1 to about 5 carbon atoms, lower alkenyl of from 2 to about 5 carbon atoms, lower alkoxy of from 1 to about 5 carbons atoms, fluoro, chloro, acetamido, amino, nitro, lower alkylamino of from 1 to about 5 carbon atoms,
10 hydroxy, lower hydroxyalkyl of from 1 to about 5 carbon atoms, cyano, or a group of the formula



- 15 wherein n is an integer from 0 to about 5.

5. The compound of claim 4, wherein Y is a straight or branched carbon chain of from 1 to about 4 carbon atoms and R is lower
20 alkyl of from 1 to about 4 carbon atoms aryl of from 6 to about 8 carbon atoms or aralkyl, wherein the alkyl portion contains from 1 to about 4 carbon atoms and the aryl portion contains from 6 to about 8 carbon atoms, and x is 1 or 2.

- 25 6. The compound of claim 5 wherein Ar is unsubstituted phenyl or phenyl substituted with lower alkyl of from 1 to about 5 carbon atoms, fluoro, chloro, nitro or a group of the formula




wherein n is an integer from 1 to about 5, and x is 1.

7. The compound of claim 5 wherein Ar is 2-methylphenyl, and x
35 is 1.

8. The compound of claim 7 wherein R is methyl or ethyl.


9. The compound of the formula



10 wherein R is methyl or ethyl, or a pharmaceutically acceptable salt thereof.

10. The compound of claim 1, 2, 4, 5, 7, 8 or 9 as the hydrochloride, sulfate, phosphate, gluconate or tartrate acid addition salt.

11. A method for the treatment or prophylaxis of cardiac disorders in a mammal, comprising administering to such mammal a short acting β -blocker of the formula



wherein Y is a straight or branched carbon chain of from 1 to about 10 carbon atoms or aralkyl of from 8 to about 20 carbon atoms; R is lower alkyl, lower alkenyl lower alkynyl, aryl or aralkyl; x is an integer from 1 to about 3, provided that when x is greater than 1, different occurrences of the -COOR group may be the same or different; Ar is unsubstituted aromatic or aromatic substituted with lower alkyl, lower alkenyl, lower alkynyl, lower alkoxy, halogen, acetamido, amino, nitro, lower alkylamino, hydroxy, lower hydroxyalkyl, cyano, or a group of the formula



0053434

wherein n is an integer from 0 to about 10; or a pharmaceutically acceptable salt thereof.


12. The method of claim 11 wherein R is lower alkyl of from 1 to about 10 carbon atoms, lower alkenyl of from 2 to about 10 carbon atoms, lower alkynyl of from 2 to about 10 carbon atoms, aryl of from 6 to about 10 carbon atoms, or aralkyl, wherein the alkyl portion contains from 1 to about 10 carbon atoms and the aryl portion contains from 6 to about 10 carbon atoms.

10

13. The method of claim 11, wherein Y is a straight or branched carbon chain of from 1 to about 6 carbon atoms or aralkyl of from 8 to about 12 carbon atoms, and R is lower alkyl of from 1 to about 5 carbon atoms, lower alkenyl of from 2 to about 5 carbon atoms, lower alkynyl of from 2 to about 5 carbon atoms, or aralkyl wherein the alkyl portion contains from 1 to about 5 carbon atoms and the aryl portion contains from 6 to about 10 carbon atoms.

14. The method of claim 13 wherein Ar is unsubstituted aromatic or aromatic substituted with lower alkyl of from 1 to about 5 carbon atoms, lower alkenyl of from 2 to about 5 carbon atoms, lower alkynyl of from 2 to about 5 carbon atoms, lower alkoxy of from 1 to about 5 carbon atoms, fluoro, chloro, acetamido, amino, nitro, lower alkylamino of from 1 to about 5 carbon atoms, hydroxy, lower hydroxyalkyl of from 1 to about 5 carbon atoms, cyano, or a group of the formula

25




30 wherein n is an integer from 0 to about 5.

15. The method of Claim 14, wherein Y is a straight or branched carbon chain of from 1 to about 4 carbon atoms and R is lower alkyl of from 1 to about 4 carbon atoms, aryl, of from 6 to about 8 carbon atoms or aralkyl, wherein the alkyl portions contain from 1 to about 4 carbon atoms and the aryl portion contains from 6 to about 8 carbon atoms, or aralkyl, wherein the alkyl portions contain from 1 to about 4 carbon

35

atoms and the aryl portion contains from 6 to about 8 carbon atoms, and x is 1 or 2.

16. The method of claim 15 wherein Ar is unsubstituted phenyl or phenyl substituted with lower alkyl of from 1 to about 5 carbon atoms, fluoro, chloro, nitro or a group of the formula



10


wherein n is an integer from 1 to about 5, and x is 1.

17. The method of Claim 15 wherein Ar is 2-methylphenyl.

18. The method of claim 17 wherein R is methyl or ethyl.

19. The method of claim 11 wherein the compound is of the formula

20



- 25 wherein R is methyl or ethyl, or a pharmaceutically acceptable salt thereof.

20. The method of claim 11, 12, 14, 15, 17, 18 or 19 wherein said compound is the hydrochloride, sulfate, phosphate, gluconate or tartrate acid addition salt.

21. The method of claim 11, 12, 14, 15, 17, 18 or 19 wherein the compound is administered parenterally.

22. the method of claim 21 wherein the compound is administered by intravenous injection or intravenous infusion at a dosage rate of from about 0.001 to about 100 mg of compound per kg of body weight of

0053434

-33-

said mammal per mg of compound per kg of body weight of said mammal per hour.

23. The method of claim 21 wherein the compound is administered
5 by intravenous injection or intravenous infusion at a dosage rate of
from about 0.01 to about 10 mg of compound per kg of body weight of said
mammal per hour.

10



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT
which under Rule 45 of the European Patent Convention
shall be considered, for the purposes of subsequent
proceedings, as the European search report

0053434

Application number

EP 81 30 5128.1

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int. Cl.)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
X	DE - A - 1 922 003 (BOEHRINGER) * examples * & GB - A - 1 245 219 -----	1-21	C 07 C 101/18 C 07 C 101/62 C 07 C 101/00
			TECHNICAL FIELDS SEARCHED (Int. Cl.)
			C 07 C 101/00
INCOMPLETE SEARCH			CATEGORY OF CITED DOCUMENTS
<p>The Search Division considers that the present European patent application does not comply with the provisions of the European Patent Convention to such an extent that it is not possible to carry out a meaningful search into the state of the art on the basis of some of the claims.</p> <p>Claims searched completely: 1-21 Claims searched incompletely: Claims not searched: 22-23 Reason for the limitation of the search:</p> <p>Method for treatment of the animal or human body by therapy (see article 52 (4) of the European Patent Convention).</p>			<p>X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document T: theory or principle underlying the invention E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons</p> <p>&: member of the same patent family, corresponding document</p>
Place of search	Berlin	Date of completion of the search	22-02-1982
		Examiner	BREW